


Age-Adjusted Invasive Cancer Incidence Rates for the 10 Primary Sites with the Highest Rates within State- and Sex-Specific Categories




Alabama,^{*†} All Races, 2001

Males



Females



* Data are from selected statewide and metropolitan area cancer registries that met the data quality criteria for publication in *United States Cancer Statistics: 2001 Incidence and Mortality*: case ascertainment is at least 90% complete; $\geq 97\%$ of cases pass a standard set of computerized edits; $\leq 5\%$ of cases were ascertained by death certificate only; $\leq 3\%$ of cases are missing information on sex; $\leq 5\%$ of cases are missing information on race; $\leq 3\%$ of cases are missing information on age; states that are included are shaded in the U.S. map above. U.S. rates cover approximately 92% of the population.

[†] Excludes basal and squamous cell carcinomas of the skin except when these occur on the skin of the genital organs, and *in situ* cancers except urinary bladder.
[‡] Rates are age-adjusted to the 2000 U.S. standard population. Rates are suppressed and not ranked if case counts are fewer than 16 or if the population of the specific sex category is less than 50,000.


Note: ONS - Other Nervous System; NOS - Not Otherwise Specified; IBD - Intrahepatic Bile Duct. Urinary bladder cases include invasive and *in situ*.

Age-Adjusted Cancer Death Rates for the 10 Primary Sites with the Highest Rates within State- and Sex-Specific Categories




Alabama,* All Races, 2001

Males



Females



* Data are from the National Vital Statistics System (NVSS).

† Rates are age-adjusted to the 2000 U.S. standard population. Rates are suppressed and not ranked if case counts are fewer than 16 or if the population of the specific sex category is less than 50,000.

Note: ONS - Other Nervous System; NOS - Not Otherwise Specified; IBD - Intrahepatic Bile Duct.



Age-Adjusted Invasive Cancer Incidence Rates for the 10 Primary Sites with the Highest Rates within State- and Sex-Specific Categories



Alaska,^{*†} All Races, 2001

Males

Females



* Data are from selected statewide and metropolitan area cancer registries that met the data quality criteria for publication in *United States Cancer Statistics: 2001 Incidence and Mortality*: case ascertainment is at least 90% complete; $\geq 97\%$ of cases pass a standard set of computerized edits; $\leq 5\%$ of cases were ascertained by death certificate only; $\leq 3\%$ of cases are missing information on sex; $\leq 5\%$ of cases are missing information on race; $\leq 3\%$ of cases are missing information on age; states that are included are shaded in the U.S. map above. U.S. rates cover approximately 92% of the population.



[†] Excludes basal and squamous cell carcinomas of the skin except when these occur on the skin of the genital organs, and *in situ* cancers except urinary bladder.
[‡] Rates are age-adjusted to the 2000 U.S. standard population. Rates are suppressed and not ranked if case counts are fewer than 16 or if the population of the specific sex category is less than 50,000.

Note: ONS - Other Nervous System; NOS - Not Otherwise Specified; IBD - Intrahepatic Bile Duct. Urinary bladder cases include invasive and *in situ*.

Age-Adjusted Cancer Death Rates for the 10 Primary Sites with the Highest Rates within State- and Sex-Specific Categories



Alaska,* All Races, 2001



* Data are from the National Vital Statistics System (NVSS).

† Rates are age-adjusted to the 2000 U.S. standard population. Rates are suppressed and not ranked if case counts are fewer than 16 or if the population of the specific sex category is less than 50,000.


Note: ONS - Other Nervous System; NOS - Not Otherwise Specified; IBD - Intrahepatic Bile Duct.

Age-Adjusted Invasive Cancer Incidence Rates for the 10 Primary Sites with the Highest Rates within State- and Sex-Specific Categories




Arizona,^{*†} All Races, 2001

Males



Females



* Data are from selected statewide and metropolitan area cancer registries that met the data quality criteria for publication in *United States Cancer Statistics: 2001 Incidence and Mortality*: case ascertainment is at least 90% complete; $\geq 97\%$ of cases pass a standard set of computerized edits; $\leq 5\%$ of cases were ascertained by death certificate only; $\leq 3\%$ of cases are missing information on sex; $\leq 5\%$ of cases are missing information on race; $\leq 3\%$ of cases are missing information on age; states that are included are shaded in the U.S. map above. U.S. rates cover approximately 92% of the population.

[†] Excludes basal and squamous cell carcinomas of the skin except when these occur on the skin of the genital organs, and *in situ* cancers except urinary bladder.
[‡] Rates are age-adjusted to the 2000 U.S. standard population. Rates are suppressed and not ranked if case counts are fewer than 16 or if the population of the specific sex category is less than 50,000.

Note: ONS - Other Nervous System; NOS - Not Otherwise Specified; IBD - Intrahepatic Bile Duct. Urinary bladder cases include invasive and *in situ*.



Age-Adjusted Cancer Death Rates for the 10 Primary Sites with the Highest Rates within State- and Sex-Specific Categories



Arizona,* All Races, 2001

Males

Females



* Data are from the National Vital Statistics System (NVSS).

† Rates are age-adjusted to the 2000 U.S. standard population. Rates are suppressed and not ranked if case counts are fewer than 16 or if the population of the specific sex category is less than 50,000.

Note: ONS - Other Nervous System; NOS - Not Otherwise Specified; IBD - Intrahepatic Bile Duct.



Age-Adjusted Invasive Cancer Incidence Rates for the 10 Primary Sites with the Highest Rates within State- and Sex-Specific Categories



Arkansas,^{*†} All Races, 2001

Males

Females



* Data are from selected statewide and metropolitan area cancer registries that met the data quality criteria for publication in *United States Cancer Statistics: 2001 Incidence and Mortality*: case ascertainment is at least 90% complete; $\geq 97\%$ of cases pass a standard set of computerized edits; $\leq 5\%$ of cases were ascertained by death certificate only; $\leq 3\%$ of cases are missing information on sex; $\leq 5\%$ of cases are missing information on race; $\leq 3\%$ of cases are missing information on age; states that are included are shaded in the U.S. map above. U.S. rates cover approximately 92% of the population.

† Excludes basal and squamous cell carcinomas of the skin except when these occur on the skin of the genital organs, and *in situ* cancers except urinary bladder.

‡ Rates are age-adjusted to the 2000 U.S. standard population. Rates are suppressed and not ranked if case counts are fewer than 16 or if the population of the specific sex category is less than 50,000.


Note: ONS - Other Nervous System; NOS - Not Otherwise Specified; IBD - Intrahepatic Bile Duct. Urinary bladder cases include invasive and *in situ*.

Age-Adjusted Cancer Death Rates for the 10 Primary Sites with the Highest Rates within State- and Sex-Specific Categories




Arkansas,* All Races, 2001

Males



Females



* Data are from the National Vital Statistics System (NVSS).

† Rates are age-adjusted to the 2000 U.S. standard population. Rates are suppressed and not ranked if case counts are fewer than 16 or if the population of the specific sex category is less than 50,000.


Note: ONS - Other Nervous System; NOS - Not Otherwise Specified; IBD - Intrahepatic Bile Duct.

Age-Adjusted Invasive Cancer Incidence Rates for the 10 Primary Sites with the Highest Rates within State- and Sex-Specific Categories




California,^{*,†} All Races, 2001

Males



Females



* Data are from selected statewide and metropolitan area cancer registries that met the data quality criteria for publication in *United States Cancer Statistics: 2001 Incidence and Mortality*: case ascertainment is at least 90% complete; $\geq 97\%$ of cases pass a standard set of computerized edits; $\leq 5\%$ of cases were ascertained by death certificate only; $\leq 3\%$ of cases are missing information on sex; $\leq 5\%$ of cases are missing information on race; $\leq 3\%$ of cases are missing information on age; states that are included are shaded in the U.S. map above. U.S. rates cover approximately 92% of the population.

[†] Excludes basal and squamous cell carcinomas of the skin except when these occur on the skin of the genital organs, and *in situ* cancers except urinary bladder.
[‡] Rates are age-adjusted to the 2000 U.S. standard population. Rates are suppressed and not ranked if case counts are fewer than 16 or if the population of the specific sex category is less than 50,000.


Note: ONS - Other Nervous System; NOS - Not Otherwise Specified; IBD - Intrahepatic Bile Duct. Urinary bladder cases include invasive and *in situ*.

Age-Adjusted Cancer Death Rates for the 10 Primary Sites with the Highest Rates within State- and Sex-Specific Categories




California,* All Races, 2001

Males



Females



* Data are from the National Vital Statistics System (NVSS).

† Rates are age-adjusted to the 2000 U.S. standard population. Rates are suppressed and not ranked if case counts are fewer than 16 or if the population of the specific sex category is less than 50,000.

Note: ONS - Other Nervous System; NOS - Not Otherwise Specified; IBD - Intrahepatic Bile Duct.



Age-Adjusted Invasive Cancer Incidence Rates for the 10 Primary Sites with the Highest Rates within State- and Sex-Specific Categories



Colorado,^{*†} All Races, 2001

Males

Females



* Data are from selected statewide and metropolitan area cancer registries that met the data quality criteria for publication in *United States Cancer Statistics: 2001 Incidence and Mortality*: case ascertainment is at least 90% complete; $\geq 97\%$ of cases pass a standard set of computerized edits; $\leq 5\%$ of cases were ascertained by death certificate only; $\leq 3\%$ of cases are missing information on sex; $\leq 5\%$ of cases are missing information on race; $\leq 3\%$ of cases are missing information on age; states that are included are shaded in the U.S. map above. U.S. rates cover approximately 92% of the population.

[†] Excludes basal and squamous cell carcinomas of the skin except when these occur on the skin of the genital organs, and *in situ* cancers except urinary bladder.
[‡] Rates are age-adjusted to the 2000 U.S. standard population. Rates are suppressed and not ranked if case counts are fewer than 16 or if the population of the specific sex category is less than 50,000.

Note: ONS - Other Nervous System; NOS - Not Otherwise Specified; IBD - Intrahepatic Bile Duct. Urinary bladder cases include invasive and *in situ*.



Age-Adjusted Cancer Death Rates for the 10 Primary Sites with the Highest Rates within State- and Sex-Specific Categories



Colorado,* All Races, 2001

Males

Females



* Data are from the National Vital Statistics System (NVSS).

† Rates are age-adjusted to the 2000 U.S. standard population. Rates are suppressed and not ranked if case counts are fewer than 16 or if the population of the specific sex category is less than 50,000.

Note: ONS - Other Nervous System; NOS - Not Otherwise Specified; IBD - Intrahepatic Bile Duct.



Age-Adjusted Invasive Cancer Incidence Rates for the 10 Primary Sites with the Highest Rates within State- and Sex-Specific Categories



Connecticut,^{*†} All Races, 2001

Males

Females



* Data are from selected statewide and metropolitan area cancer registries that met the data quality criteria for publication in *United States Cancer Statistics: 2001 Incidence and Mortality*: case ascertainment is at least 90% complete; $\geq 97\%$ of cases pass a standard set of computerized edits; $\leq 5\%$ of cases were ascertained by death certificate only; $\leq 3\%$ of cases are missing information on sex; $\leq 5\%$ of cases are missing information on race; $\leq 3\%$ of cases are missing information on age; states that are included are shaded in the U.S. map above. U.S. rates cover approximately 92% of the population.

[†] Excludes basal and squamous cell carcinomas of the skin except when these occur on the skin of the genital organs, and *in situ* cancers except urinary bladder.
[‡] Rates are age-adjusted to the 2000 U.S. standard population. Rates are suppressed and not ranked if case counts are fewer than 16 or if the population of the specific sex category is less than 50,000.

Note: ONS - Other Nervous System; NOS - Not Otherwise Specified; IBD - Intrahepatic Bile Duct. Urinary bladder cases include invasive and *in situ*.



Age-Adjusted Cancer Death Rates for the 10 Primary Sites with the Highest Rates within State- and Sex-Specific Categories



Connecticut,* All Races, 2001

Males

Females



* Data are from the National Vital Statistics System (NVSS).

† Rates are age-adjusted to the 2000 U.S. standard population. Rates are suppressed and not ranked if case counts are fewer than 16 or if the population of the specific sex category is less than 50,000.

Note: ONS - Other Nervous System; NOS - Not Otherwise Specified; IBD - Intrahepatic Bile Duct.



Age-Adjusted Cancer Death Rates for the 10 Primary Sites with the Highest Rates within State- and Sex-Specific Categories



Delaware,* All Races, 2001

Males

Females



* Data are from the National Vital Statistics System (NVSS).

† Rates are age-adjusted to the 2000 U.S. standard population. Rates are suppressed and not ranked if case counts are fewer than 16 or if the population of the specific sex category is less than 50,000.

Note: ONS - Other Nervous System; NOS - Not Otherwise Specified; IBD - Intrahepatic Bile Duct.



Age-Adjusted Invasive Cancer Incidence Rates for the 10 Primary Sites with the Highest Rates within State- and Sex-Specific Categories



District of Columbia,^{*†} All Races, 2001

Males

Females



* Data are from selected statewide and metropolitan area cancer registries that met the data quality criteria for publication in *United States Cancer Statistics: 2001 Incidence and Mortality*: case ascertainment is at least 90% complete; $\geq 97\%$ of cases pass a standard set of computerized edits; $\leq 5\%$ of cases were ascertained by death certificate only; $\leq 3\%$ of cases are missing information on sex; $\leq 5\%$ of cases are missing information on race; $\leq 3\%$ of cases are missing information on age; states that are included are shaded in the U.S. map above. U.S. rates cover approximately 92% of the population.

[†] Excludes basal and squamous cell carcinomas of the skin except when these occur on the skin of the genital organs, and *in situ* cancers except urinary bladder.
[‡] Rates are age-adjusted to the 2000 U.S. standard population. Rates are suppressed and not ranked if case counts are fewer than 16 or if the population of the specific sex category is less than 50,000.


Note: ONS - Other Nervous System; NOS - Not Otherwise Specified; IBD - Intrahepatic Bile Duct. Urinary bladder cases include invasive and *in situ*.

Age-Adjusted Cancer Death Rates for the 10 Primary Sites with the Highest Rates within State- and Sex-Specific Categories




District of Columbia,* All Races, 2001

Males



Females



* Data are from the National Vital Statistics System (NVSS).

† Rates are age-adjusted to the 2000 U.S. standard population. Rates are suppressed and not ranked if case counts are fewer than 16 or if the population of the specific sex category is less than 50,000.


Note: ONS - Other Nervous System; NOS - Not Otherwise Specified; IBD - Intrahepatic Bile Duct.

Age-Adjusted Invasive Cancer Incidence Rates for the 10 Primary Sites with the Highest Rates within State- and Sex-Specific Categories




Florida,^{*†} All Races, 2001

Males



Females



* Data are from selected statewide and metropolitan area cancer registries that met the data quality criteria for publication in *United States Cancer Statistics: 2001 Incidence and Mortality*: case ascertainment is at least 90% complete; $\geq 97\%$ of cases pass a standard set of computerized edits; $\leq 5\%$ of cases were ascertained by death certificate only; $\leq 3\%$ of cases are missing information on sex; $\leq 5\%$ of cases are missing information on race; $\leq 3\%$ of cases are missing information on age; states that are included are shaded in the U.S. map above. U.S. rates cover approximately 92% of the population.

[†] Excludes basal and squamous cell carcinomas of the skin except when these occur on the skin of the genital organs, and *in situ* cancers except urinary bladder.
[‡] Rates are age-adjusted to the 2000 U.S. standard population. Rates are suppressed and not ranked if case counts are fewer than 16 or if the population of the specific sex category is less than 50,000.


Note: ONS - Other Nervous System; NOS - Not Otherwise Specified; IBD - Intrahepatic Bile Duct. Urinary bladder cases include invasive and *in situ*.

Age-Adjusted Cancer Death Rates for the 10 Primary Sites with the Highest Rates within State- and Sex-Specific Categories




Florida,* All Races, 2001

Males



Females



* Data are from the National Vital Statistics System (NVSS).

† Rates are age-adjusted to the 2000 U.S. standard population. Rates are suppressed and not ranked if case counts are fewer than 16 or if the population of the specific sex category is less than 50,000.


Note: ONS - Other Nervous System; NOS - Not Otherwise Specified; IBD - Intrahepatic Bile Duct.

Age-Adjusted Invasive Cancer Incidence Rates for the 10 Primary Sites with the Highest Rates within State- and Sex-Specific Categories




Georgia,^{*†} All Races, 2001

Males



Females



* Data are from selected statewide and metropolitan area cancer registries that met the data quality criteria for publication in *United States Cancer Statistics: 2001 Incidence and Mortality*: case ascertainment is at least 90% complete; $\geq 97\%$ of cases pass a standard set of computerized edits; $\leq 5\%$ of cases were ascertained by death certificate only; $\leq 3\%$ of cases are missing information on sex; $\leq 5\%$ of cases are missing information on race; $\leq 3\%$ of cases are missing information on age; states that are included are shaded in the U.S. map above. U.S. rates cover approximately 92% of the population.

[†] Excludes basal and squamous cell carcinomas of the skin except when these occur on the skin of the genital organs, and *in situ* cancers except urinary bladder.

[‡] Rates are age-adjusted to the 2000 U.S. standard population. Rates are suppressed and not ranked if case counts are fewer than 16 or if the population of the specific sex category is less than 50,000.


Note: ONS - Other Nervous System; NOS - Not Otherwise Specified; IBD - Intrahepatic Bile Duct. Urinary bladder cases include invasive and *in situ*.

Age-Adjusted Cancer Death Rates for the 10 Primary Sites with the Highest Rates within State- and Sex-Specific Categories




Georgia,* All Races, 2001

Males



Females



* Data are from the National Vital Statistics System (NVSS).

† Rates are age-adjusted to the 2000 U.S. standard population. Rates are suppressed and not ranked if case counts are fewer than 16 or if the population of the specific sex category is less than 50,000.


Note: ONS - Other Nervous System; NOS - Not Otherwise Specified; IBD - Intrahepatic Bile Duct.

Age-Adjusted Invasive Cancer Incidence Rates for the 10 Primary Sites with the Highest Rates within State- and Sex-Specific Categories




Hawaii,^{*†} All Races, 2001

Males



Females



* Data are from selected statewide and metropolitan area cancer registries that met the data quality criteria for publication in *United States Cancer Statistics: 2001 Incidence and Mortality*: case ascertainment is at least 90% complete; $\geq 97\%$ of cases pass a standard set of computerized edits; $\leq 5\%$ of cases were ascertained by death certificate only; $\leq 3\%$ of cases are missing information on sex; $\leq 5\%$ of cases are missing information on race; $\leq 3\%$ of cases are missing information on age; states that are included are shaded in the U.S. map above. U.S. rates cover approximately 92% of the population.

[†] Excludes basal and squamous cell carcinomas of the skin except when these occur on the skin of the genital organs, and *in situ* cancers except urinary bladder.

[‡] Rates are age-adjusted to the 2000 U.S. standard population. Rates are suppressed and not ranked if case counts are fewer than 16 or if the population of the specific sex category is less than 50,000.


Note: ONS - Other Nervous System; NOS - Not Otherwise Specified; IBD - Intrahepatic Bile Duct. Urinary bladder cases include invasive and *in situ*.

Age-Adjusted Cancer Death Rates for the 10 Primary Sites with the Highest Rates within State- and Sex-Specific Categories




Hawaii,* All Races, 2001

Males



Females



* Data are from the National Vital Statistics System (NVSS).

† Rates are age-adjusted to the 2000 U.S. standard population. Rates are suppressed and not ranked if case counts are fewer than 16 or if the population of the specific sex category is less than 50,000.

Note: ONS - Other Nervous System; NOS - Not Otherwise Specified; IBD - Intrahepatic Bile Duct.



Age-Adjusted Invasive Cancer Incidence Rates for the 10 Primary Sites with the Highest Rates within State- and Sex-Specific Categories



Idaho,^{*†} All Races, 2001

Males

Females



* Data are from selected statewide and metropolitan area cancer registries that met the data quality criteria for publication in *United States Cancer Statistics: 2001 Incidence and Mortality*: case ascertainment is at least 90% complete; $\geq 97\%$ of cases pass a standard set of computerized edits; $\leq 5\%$ of cases were ascertained by death certificate only; $\leq 3\%$ of cases are missing information on sex; $\leq 5\%$ of cases are missing information on race; $\leq 3\%$ of cases are missing information on age; states that are included are shaded in the U.S. map above. U.S. rates cover approximately 92% of the population.

† Excludes basal and squamous cell carcinomas of the skin except when these occur on the skin of the genital organs, and *in situ* cancers except urinary bladder.

‡ Rates are age-adjusted to the 2000 U.S. standard population. Rates are suppressed and not ranked if case counts are fewer than 16 or if the population of the specific sex category is less than 50,000.

Note: ONS - Other Nervous System; NOS - Not Otherwise Specified; IBD - Intrahepatic Bile Duct. Urinary bladder cases include invasive and *in situ*.



Age-Adjusted Cancer Death Rates for the 10 Primary Sites with the Highest Rates within State- and Sex-Specific Categories



Idaho,* All Races, 2001

Males

Females



* Data are from the National Vital Statistics System (NVSS).

† Rates are age-adjusted to the 2000 U.S. standard population. Rates are suppressed and not ranked if case counts are fewer than 16 or if the population of the specific sex category is less than 50,000.

Note: ONS - Other Nervous System; NOS - Not Otherwise Specified; IBD - Intrahepatic Bile Duct.



Age-Adjusted Invasive Cancer Incidence Rates for the 10 Primary Sites with the Highest Rates within State- and Sex-Specific Categories



Illinois,^{*†} All Races, 2001

Males

Females



* Data are from selected statewide and metropolitan area cancer registries that met the data quality criteria for publication in *United States Cancer Statistics: 2001 Incidence and Mortality*: case ascertainment is at least 90% complete; $\geq 97\%$ of cases pass a standard set of computerized edits; $\leq 5\%$ of cases were ascertained by death certificate only; $\leq 3\%$ of cases are missing information on sex; $\leq 5\%$ of cases are missing information on race; $\leq 3\%$ of cases are missing information on age; states that are included are shaded in the U.S. map above. U.S. rates cover approximately 92% of the population.

[†] Excludes basal and squamous cell carcinomas of the skin except when these occur on the skin of the genital organs, and *in situ* cancers except urinary bladder.

[‡] Rates are age-adjusted to the 2000 U.S. standard population. Rates are suppressed and not ranked if case counts are fewer than 16 or if the population of the specific sex category is less than 50,000.

Note: ONS - Other Nervous System; NOS - Not Otherwise Specified; IBD - Intrahepatic Bile Duct. Urinary bladder cases include invasive and *in situ*.



Age-Adjusted Cancer Death Rates for the 10 Primary Sites with the Highest Rates within State- and Sex-Specific Categories



Illinois,* All Races, 2001

Males

Females



* Data are from the National Vital Statistics System (NVSS).

† Rates are age-adjusted to the 2000 U.S. standard population. Rates are suppressed and not ranked if case counts are fewer than 16 or if the population of the specific sex category is less than 50,000.

Note: ONS - Other Nervous System; NOS - Not Otherwise Specified; IBD - Intrahepatic Bile Duct.



Age-Adjusted Invasive Cancer Incidence Rates for the 10 Primary Sites with the Highest Rates within State- and Sex-Specific Categories



Indiana,^{*,†} All Races, 2001

Males

Females



* Data are from selected statewide and metropolitan area cancer registries that met the data quality criteria for publication in *United States Cancer Statistics: 2001 Incidence and Mortality*: case ascertainment is at least 90% complete; $\geq 97\%$ of cases pass a standard set of computerized edits; $\leq 5\%$ of cases were ascertained by death certificate only; $\leq 3\%$ of cases are missing information on sex; $\leq 5\%$ of cases are missing information on race; $\leq 3\%$ of cases are missing information on age; states that are included are shaded in the U.S. map above. U.S. rates cover approximately 92% of the population.

[†] Excludes basal and squamous cell carcinomas of the skin except when these occur on the skin of the genital organs, and *in situ* cancers except urinary bladder.
[‡] Rates are age-adjusted to the 2000 U.S. standard population. Rates are suppressed and not ranked if case counts are fewer than 16 or if the population of the specific sex category is less than 50,000.


Note: ONS - Other Nervous System; NOS - Not Otherwise Specified; IBD - Intrahepatic Bile Duct. Urinary bladder cases include invasive and *in situ*.

Age-Adjusted Cancer Death Rates for the 10 Primary Sites with the Highest Rates within State- and Sex-Specific Categories




Indiana,* All Races, 2001

Males



Females



* Data are from the National Vital Statistics System (NVSS).

† Rates are age-adjusted to the 2000 U.S. standard population. Rates are suppressed and not ranked if case counts are fewer than 16 or if the population of the specific sex category is less than 50,000.

Note: ONS - Other Nervous System; NOS - Not Otherwise Specified; IBD - Intrahepatic Bile Duct.



Age-Adjusted Invasive Cancer Incidence Rates for the 10 Primary Sites with the Highest Rates within State- and Sex-Specific Categories



Iowa,^{*†} All Races, 2001

Males

Females



* Data are from selected statewide and metropolitan area cancer registries that met the data quality criteria for publication in *United States Cancer Statistics: 2001 Incidence and Mortality*: case ascertainment is at least 90% complete; $\geq 97\%$ of cases pass a standard set of computerized edits; $\leq 5\%$ of cases were ascertained by death certificate only; $\leq 3\%$ of cases are missing information on sex; $\leq 5\%$ of cases are missing information on race; $\leq 3\%$ of cases are missing information on age; states that are included are shaded in the U.S. map above. U.S. rates cover approximately 92% of the population.

[†] Excludes basal and squamous cell carcinomas of the skin except when these occur on the skin of the genital organs, and *in situ* cancers except urinary bladder.

[‡] Rates are age-adjusted to the 2000 U.S. standard population. Rates are suppressed and not ranked if case counts are fewer than 16 or if the population of the specific sex category is less than 50,000.

Note: ONS - Other Nervous System; NOS - Not Otherwise Specified; IBD - Intrahepatic Bile Duct. Urinary bladder cases include invasive and *in situ*.



Age-Adjusted Cancer Death Rates for the 10 Primary Sites with the Highest Rates within State- and Sex-Specific Categories



Iowa,* All Races, 2001

Males

Females



* Data are from the National Vital Statistics System (NVSS).

† Rates are age-adjusted to the 2000 U.S. standard population. Rates are suppressed and not ranked if case counts are fewer than 16 or if the population of the specific sex category is less than 50,000.

Note: ONS - Other Nervous System; NOS - Not Otherwise Specified; IBD - Intrahepatic Bile Duct.



Age-Adjusted Invasive Cancer Incidence Rates for the 10 Primary Sites with the Highest Rates within State- and Sex-Specific Categories



Kansas,^{*†} All Races, 2001

Males

Females



* Data are from selected statewide and metropolitan area cancer registries that met the data quality criteria for publication in *United States Cancer Statistics: 2001 Incidence and Mortality*: case ascertainment is at least 90% complete; $\geq 97\%$ of cases pass a standard set of computerized edits; $\leq 5\%$ of cases were ascertained by death certificate only; $\leq 3\%$ of cases are missing information on sex; $\leq 5\%$ of cases are missing information on race; $\leq 3\%$ of cases are missing information on age; states that are included are shaded in the U.S. map above. U.S. rates cover approximately 92% of the population.

[†] Excludes basal and squamous cell carcinomas of the skin except when these occur on the skin of the genital organs, and *in situ* cancers except urinary bladder.
[‡] Rates are age-adjusted to the 2000 U.S. standard population. Rates are suppressed and not ranked if case counts are fewer than 16 or if the population of the specific sex category is less than 50,000.

Note: ONS - Other Nervous System; NOS - Not Otherwise Specified; IBD - Intrahepatic Bile Duct. Urinary bladder cases include invasive and *in situ*.



Age-Adjusted Cancer Death Rates for the 10 Primary Sites with the Highest Rates within State- and Sex-Specific Categories



Kansas,* All Races, 2001

Males

Females



* Data are from the National Vital Statistics System (NVSS).

† Rates are age-adjusted to the 2000 U.S. standard population. Rates are suppressed and not ranked if case counts are fewer than 16 or if the population of the specific sex category is less than 50,000.

Note: ONS - Other Nervous System; NOS - Not Otherwise Specified; IBD - Intrahepatic Bile Duct.



Age-Adjusted Invasive Cancer Incidence Rates for the 10 Primary Sites with the Highest Rates within State- and Sex-Specific Categories



Kentucky,^{*,†} All Races, 2001

Males

Females



* Data are from selected statewide and metropolitan area cancer registries that met the data quality criteria for publication in *United States Cancer Statistics: 2001 Incidence and Mortality*: case ascertainment is at least 90% complete; $\geq 97\%$ of cases pass a standard set of computerized edits; $\leq 5\%$ of cases were ascertained by death certificate only; $\leq 3\%$ of cases are missing information on sex; $\leq 5\%$ of cases are missing information on race; $\leq 3\%$ of cases are missing information on age; states that are included are shaded in the U.S. map above. U.S. rates cover approximately 92% of the population.

[†] Excludes basal and squamous cell carcinomas of the skin except when these occur on the skin of the genital organs, and *in situ* cancers except urinary bladder.
[‡] Rates are age-adjusted to the 2000 U.S. standard population. Rates are suppressed and not ranked if case counts are fewer than 16 or if the population of the specific sex category is less than 50,000.

Note: ONS - Other Nervous System; NOS - Not Otherwise Specified; IBD - Intrahepatic Bile Duct. Urinary bladder cases include invasive and *in situ*.



Age-Adjusted Cancer Death Rates for the 10 Primary Sites with the Highest Rates within State- and Sex-Specific Categories



Kentucky,* All Races, 2001

Males

Females



* Data are from the National Vital Statistics System (NVSS).

† Rates are age-adjusted to the 2000 U.S. standard population. Rates are suppressed and not ranked if case counts are fewer than 16 or if the population of the specific sex category is less than 50,000.

Note: ONS - Other Nervous System; NOS - Not Otherwise Specified; IBD - Intrahepatic Bile Duct.



Age-Adjusted Invasive Cancer Incidence Rates for the 10 Primary Sites with the Highest Rates within State- and Sex-Specific Categories



Louisiana,^{*†} All Races, 2001

Males

Females



* Data are from selected statewide and metropolitan area cancer registries that met the data quality criteria for publication in *United States Cancer Statistics: 2001 Incidence and Mortality*: case ascertainment is at least 90% complete; $\geq 97\%$ of cases pass a standard set of computerized edits; $\leq 5\%$ of cases were ascertained by death certificate only; $\leq 3\%$ of cases are missing information on sex; $\leq 5\%$ of cases are missing information on race; $\leq 3\%$ of cases are missing information on age; states that are included are shaded in the U.S. map above. U.S. rates cover approximately 92% of the population.

[†] Excludes basal and squamous cell carcinomas of the skin except when these occur on the skin of the genital organs, and *in situ* cancers except urinary bladder.
[‡] Rates are age-adjusted to the 2000 U.S. standard population. Rates are suppressed and not ranked if case counts are fewer than 16 or if the population of the specific sex category is less than 50,000.

Note: ONS - Other Nervous System; NOS - Not Otherwise Specified; IBD - Intrahepatic Bile Duct. Urinary bladder cases include invasive and *in situ*.



Age-Adjusted Cancer Death Rates for the 10 Primary Sites with the Highest Rates within State- and Sex-Specific Categories



Louisiana,* All Races, 2001

Males

Females



* Data are from the National Vital Statistics System (NVSS).

† Rates are age-adjusted to the 2000 U.S. standard population. Rates are suppressed and not ranked if case counts are fewer than 16 or if the population of the specific sex category is less than 50,000.


Note: ONS - Other Nervous System; NOS - Not Otherwise Specified; IBD - Intrahepatic Bile Duct.

Age-Adjusted Invasive Cancer Incidence Rates for the 10 Primary Sites with the Highest Rates within State- and Sex-Specific Categories




Maine,^{*}† All Races, 2001

Males



Females



* Data are from selected statewide and metropolitan area cancer registries that met the data quality criteria for publication in *United States Cancer Statistics: 2001 Incidence and Mortality*: case ascertainment is at least 90% complete; $\geq 97\%$ of cases pass a standard set of computerized edits; $\leq 5\%$ of cases were ascertained by death certificate only; $\leq 3\%$ of cases are missing information on sex; $\leq 5\%$ of cases are missing information on race; $\leq 3\%$ of cases are missing information on age; states that are included are shaded in the U.S. map above. U.S. rates cover approximately 92% of the population.

† Excludes basal and squamous cell carcinomas of the skin except when these occur on the skin of the genital organs, and *in situ* cancers except urinary bladder.
‡ Rates are age-adjusted to the 2000 U.S. standard population. Rates are suppressed and not ranked if case counts are fewer than 16 or if the population of the specific sex category is less than 50,000.

Note: ONS - Other Nervous System; NOS - Not Otherwise Specified; IBD - Intrahepatic Bile Duct. Urinary bladder cases include invasive and *in situ*.



Age-Adjusted Cancer Death Rates for the 10 Primary Sites with the Highest Rates within State- and Sex-Specific Categories



Maine,* All Races, 2001

Males

Females



* Data are from the National Vital Statistics System (NVSS).

† Rates are age-adjusted to the 2000 U.S. standard population. Rates are suppressed and not ranked if case counts are fewer than 16 or if the population of the specific sex category is less than 50,000.

Note: ONS - Other Nervous System; NOS - Not Otherwise Specified; IBD - Intrahepatic Bile Duct.



Age-Adjusted Cancer Death Rates for the 10 Primary Sites with the Highest Rates within State- and Sex-Specific Categories



Maryland,* All Races, 2001

Males

Females



* Data are from the National Vital Statistics System (NVSS).

† Rates are age-adjusted to the 2000 U.S. standard population. Rates are suppressed and not ranked if case counts are fewer than 16 or if the population of the specific sex category is less than 50,000.

Note: ONS - Other Nervous System; NOS - Not Otherwise Specified; IBD - Intrahepatic Bile Duct.



Age-Adjusted Invasive Cancer Incidence Rates for the 10 Primary Sites with the Highest Rates within State- and Sex-Specific Categories



Massachusetts,^{*†} All Races, 2001

Males

Females



* Data are from selected statewide and metropolitan area cancer registries that met the data quality criteria for publication in *United States Cancer Statistics: 2001 Incidence and Mortality*: case ascertainment is at least 90% complete; $\geq 97\%$ of cases pass a standard set of computerized edits; $\leq 5\%$ of cases were ascertained by death certificate only; $\leq 3\%$ of cases are missing information on sex; $\leq 5\%$ of cases are missing information on race; $\leq 3\%$ of cases are missing information on age; states that are included are shaded in the U.S. map above. U.S. rates cover approximately 92% of the population.

[†] Excludes basal and squamous cell carcinomas of the skin except when these occur on the skin of the genital organs, and *in situ* cancers except urinary bladder.

[‡] Rates are age-adjusted to the 2000 U.S. standard population. Rates are suppressed and not ranked if case counts are fewer than 16 or if the population of the specific sex category is less than 50,000.

Note: ONS - Other Nervous System; NOS - Not Otherwise Specified; IBD - Intrahepatic Bile Duct. Urinary bladder cases include invasive and *in situ*.



Age-Adjusted Cancer Death Rates for the 10 Primary Sites with the Highest Rates within State- and Sex-Specific Categories



Massachusetts,* All Races, 2001

Males

Females



* Data are from the National Vital Statistics System (NVSS).

† Rates are age-adjusted to the 2000 U.S. standard population. Rates are suppressed and not ranked if case counts are fewer than 16 or if the population of the specific sex category is less than 50,000.

Note: ONS - Other Nervous System; NOS - Not Otherwise Specified; IBD - Intrahepatic Bile Duct.



Age-Adjusted Invasive Cancer Incidence Rates for the 10 Primary Sites with the Highest Rates within State- and Sex-Specific Categories



Michigan,^{*†} All Races, 2001

Males

Females



* Data are from selected statewide and metropolitan area cancer registries that met the data quality criteria for publication in *United States Cancer Statistics: 2001 Incidence and Mortality*: case ascertainment is at least 90% complete; $\geq 97\%$ of cases pass a standard set of computerized edits; $\leq 5\%$ of cases were ascertained by death certificate only; $\leq 3\%$ of cases are missing information on sex; $\leq 5\%$ of cases are missing information on race; $\leq 3\%$ of cases are missing information on age; states that are included are shaded in the U.S. map above. U.S. rates cover approximately 92% of the population.

[†] Excludes basal and squamous cell carcinomas of the skin except when these occur on the skin of the genital organs, and *in situ* cancers except urinary bladder.

[‡] Rates are age-adjusted to the 2000 U.S. standard population. Rates are suppressed and not ranked if case counts are fewer than 16 or if the population of the specific sex category is less than 50,000.

Note: ONS - Other Nervous System; NOS - Not Otherwise Specified; IBD - Intrahepatic Bile Duct. Urinary bladder cases include invasive and *in situ*.



Age-Adjusted Cancer Death Rates for the 10 Primary Sites with the Highest Rates within State- and Sex-Specific Categories



Michigan,* All Races, 2001

Males

Females



* Data are from the National Vital Statistics System (NVSS).

† Rates are age-adjusted to the 2000 U.S. standard population. Rates are suppressed and not ranked if case counts are fewer than 16 or if the population of the specific sex category is less than 50,000.


Note: ONS - Other Nervous System; NOS - Not Otherwise Specified; IBD - Intrahepatic Bile Duct.

Age-Adjusted Invasive Cancer Incidence Rates for the 10 Primary Sites with the Highest Rates within State- and Sex-Specific Categories




Minnesota,^{*†} All Races, 2001

Males



Females



* Data are from selected statewide and metropolitan area cancer registries that met the data quality criteria for publication in *United States Cancer Statistics: 2001 Incidence and Mortality*: case ascertainment is at least 90% complete; $\geq 97\%$ of cases pass a standard set of computerized edits; $\leq 5\%$ of cases were ascertained by death certificate only; $\leq 3\%$ of cases are missing information on sex; $\leq 5\%$ of cases are missing information on race; $\leq 3\%$ of cases are missing information on age; states that are included are shaded in the U.S. map above. U.S. rates cover approximately 92% of the population.

[†] Excludes basal and squamous cell carcinomas of the skin except when these occur on the skin of the genital organs, and *in situ* cancers except urinary bladder.
[‡] Rates are age-adjusted to the 2000 U.S. standard population. Rates are suppressed and not ranked if case counts are fewer than 16 or if the population of the specific sex category is less than 50,000.


Note: ONS - Other Nervous System; NOS - Not Otherwise Specified; IBD - Intrahepatic Bile Duct. Urinary bladder cases include invasive and *in situ*.

Age-Adjusted Cancer Death Rates for the 10 Primary Sites with the Highest Rates within State- and Sex-Specific Categories




Minnesota,* All Races, 2001

Males



Females



* Data are from the National Vital Statistics System (NVSS).

† Rates are age-adjusted to the 2000 U.S. standard population. Rates are suppressed and not ranked if case counts are fewer than 16 or if the population of the specific sex category is less than 50,000.

Note: ONS - Other Nervous System; NOS - Not Otherwise Specified; IBD - Intrahepatic Bile Duct.



Age-Adjusted Cancer Death Rates for the 10 Primary Sites with the Highest Rates within State- and Sex-Specific Categories



Mississippi,* All Races, 2001

Males

Females



* Data are from the National Vital Statistics System (NVSS).

† Rates are age-adjusted to the 2000 U.S. standard population. Rates are suppressed and not ranked if case counts are fewer than 16 or if the population of the specific sex category is less than 50,000.

Note: ONS - Other Nervous System; NOS - Not Otherwise Specified; IBD - Intrahepatic Bile Duct.



Age-Adjusted Invasive Cancer Incidence Rates for the 10 Primary Sites with the Highest Rates within State- and Sex-Specific Categories



Missouri,^{*†} All Races, 2001

Males

Females



* Data are from selected statewide and metropolitan area cancer registries that met the data quality criteria for publication in *United States Cancer Statistics: 2001 Incidence and Mortality*: case ascertainment is at least 90% complete; $\geq 97\%$ of cases pass a standard set of computerized edits; $\leq 5\%$ of cases were ascertained by death certificate only; $\leq 3\%$ of cases are missing information on sex; $\leq 5\%$ of cases are missing information on race; $\leq 3\%$ of cases are missing information on age; states that are included are shaded in the U.S. map above. U.S. rates cover approximately 92% of the population.

[†] Excludes basal and squamous cell carcinomas of the skin except when these occur on the skin of the genital organs, and *in situ* cancers except urinary bladder.

[‡] Rates are age-adjusted to the 2000 U.S. standard population. Rates are suppressed and not ranked if case counts are fewer than 16 or if the population of the specific sex category is less than 50,000.


Note: ONS - Other Nervous System; NOS - Not Otherwise Specified; IBD - Intrahepatic Bile Duct. Urinary bladder cases include invasive and *in situ*.

Age-Adjusted Cancer Death Rates for the 10 Primary Sites with the Highest Rates within State- and Sex-Specific Categories




Missouri,* All Races, 2001

Males



Females



* Data are from the National Vital Statistics System (NVSS).

† Rates are age-adjusted to the 2000 U.S. standard population. Rates are suppressed and not ranked if case counts are fewer than 16 or if the population of the specific sex category is less than 50,000.

Note: ONS - Other Nervous System; NOS - Not Otherwise Specified; IBD - Intrahepatic Bile Duct.



Age-Adjusted Invasive Cancer Incidence Rates for the 10 Primary Sites with the Highest Rates within State- and Sex-Specific Categories



Montana,^{*†} All Races, 2001

Males

Females



* Data are from selected statewide and metropolitan area cancer registries that met the data quality criteria for publication in *United States Cancer Statistics: 2001 Incidence and Mortality*: case ascertainment is at least 90% complete; $\geq 97\%$ of cases pass a standard set of computerized edits; $\leq 5\%$ of cases were ascertained by death certificate only; $\leq 3\%$ of cases are missing information on sex; $\leq 5\%$ of cases are missing information on race; $\leq 3\%$ of cases are missing information on age; states that are included are shaded in the U.S. map above. U.S. rates cover approximately 92% of the population.

[†] Excludes basal and squamous cell carcinomas of the skin except when these occur on the skin of the genital organs, and *in situ* cancers except urinary bladder.

[‡] Rates are age-adjusted to the 2000 U.S. standard population. Rates are suppressed and not ranked if case counts are fewer than 16 or if the population of the specific sex category is less than 50,000.

Note: ONS - Other Nervous System; NOS - Not Otherwise Specified; IBD - Intrahepatic Bile Duct. Urinary bladder cases include invasive and *in situ*.



Age-Adjusted Cancer Death Rates for the 10 Primary Sites with the Highest Rates within State- and Sex-Specific Categories



Montana,* All Races, 2001

Males

Females



* Data are from the National Vital Statistics System (NVSS).

† Rates are age-adjusted to the 2000 U.S. standard population. Rates are suppressed and not ranked if case counts are fewer than 16 or if the population of the specific sex category is less than 50,000.

Note: ONS - Other Nervous System; NOS - Not Otherwise Specified; IBD - Intrahepatic Bile Duct.



Age-Adjusted Invasive Cancer Incidence Rates for the 10 Primary Sites with the Highest Rates within State- and Sex-Specific Categories



Nebraska,^{*†} All Races, 2001

Males

Females



* Data are from selected statewide and metropolitan area cancer registries that met the data quality criteria for publication in *United States Cancer Statistics: 2001 Incidence and Mortality*: case ascertainment is at least 90% complete; $\geq 97\%$ of cases pass a standard set of computerized edits; $\leq 5\%$ of cases were ascertained by death certificate only; $\leq 3\%$ of cases are missing information on sex; $\leq 5\%$ of cases are missing information on race; $\leq 3\%$ of cases are missing information on age; states that are included are shaded in the U.S. map above. U.S. rates cover approximately 92% of the population.

[†] Excludes basal and squamous cell carcinomas of the skin except when these occur on the skin of the genital organs, and *in situ* cancers except urinary bladder.
[‡] Rates are age-adjusted to the 2000 U.S. standard population. Rates are suppressed and not ranked if case counts are fewer than 16 or if the population of the specific sex category is less than 50,000.

Note: ONS - Other Nervous System; NOS - Not Otherwise Specified; IBD - Intrahepatic Bile Duct. Urinary bladder cases include invasive and *in situ*.



Age-Adjusted Cancer Death Rates for the 10 Primary Sites with the Highest Rates within State- and Sex-Specific Categories



Nebraska,* All Races, 2001

Males

Females



* Data are from the National Vital Statistics System (NVSS).

† Rates are age-adjusted to the 2000 U.S. standard population. Rates are suppressed and not ranked if case counts are fewer than 16 or if the population of the specific sex category is less than 50,000.

Note: ONS - Other Nervous System; NOS - Not Otherwise Specified; IBD - Intrahepatic Bile Duct.



Age-Adjusted Invasive Cancer Incidence Rates for the 10 Primary Sites with the Highest Rates within State- and Sex-Specific Categories



Nevada,^{*†} All Races, 2001

Males

Females



* Data are from selected statewide and metropolitan area cancer registries that met the data quality criteria for publication in *United States Cancer Statistics: 2001 Incidence and Mortality*: case ascertainment is at least 90% complete; $\geq 97\%$ of cases pass a standard set of computerized edits; $\leq 5\%$ of cases were ascertained by death certificate only; $\leq 3\%$ of cases are missing information on sex; $\leq 5\%$ of cases are missing information on race; $\leq 3\%$ of cases are missing information on age; states that are included are shaded in the U.S. map above. U.S. rates cover approximately 92% of the population.

[†] Excludes basal and squamous cell carcinomas of the skin except when these occur on the skin of the genital organs, and *in situ* cancers except urinary bladder.
[‡] Rates are age-adjusted to the 2000 U.S. standard population. Rates are suppressed and not ranked if case counts are fewer than 16 or if the population of the specific sex category is less than 50,000.


Note: ONS - Other Nervous System; NOS - Not Otherwise Specified; IBD - Intrahepatic Bile Duct. Urinary bladder cases include invasive and *in situ*.

Age-Adjusted Cancer Death Rates for the 10 Primary Sites with the Highest Rates within State- and Sex-Specific Categories




Nevada,* All Races, 2001

Males



Females



* Data are from the National Vital Statistics System (NVSS).

† Rates are age-adjusted to the 2000 U.S. standard population. Rates are suppressed and not ranked if case counts are fewer than 16 or if the population of the specific sex category is less than 50,000.


Note: ONS - Other Nervous System; NOS - Not Otherwise Specified; IBD - Intrahepatic Bile Duct.

Age-Adjusted Invasive Cancer Incidence Rates for the 10 Primary Sites with the Highest Rates within State- and Sex-Specific Categories




New Hampshire,^{*†} All Races, 2001

Males



Females



* Data are from selected statewide and metropolitan area cancer registries that met the data quality criteria for publication in *United States Cancer Statistics: 2001 Incidence and Mortality*: case ascertainment is at least 90% complete; $\geq 97\%$ of cases pass a standard set of computerized edits; $\leq 5\%$ of cases were ascertained by death certificate only; $\leq 3\%$ of cases are missing information on sex; $\leq 5\%$ of cases are missing information on race; $\leq 3\%$ of cases are missing information on age; states that are included are shaded in the U.S. map above. U.S. rates cover approximately 92% of the population.

[†] Excludes basal and squamous cell carcinomas of the skin except when these occur on the skin of the genital organs, and *in situ* cancers except urinary bladder.
[‡] Rates are age-adjusted to the 2000 U.S. standard population. Rates are suppressed and not ranked if case counts are fewer than 16 or if the population of the specific sex category is less than 50,000.


Note: ONS - Other Nervous System; NOS - Not Otherwise Specified; IBD - Intrahepatic Bile Duct. Urinary bladder cases include invasive and *in situ*.

Age-Adjusted Cancer Death Rates for the 10 Primary Sites with the Highest Rates within State- and Sex-Specific Categories




New Hampshire,* All Races, 2001

Males



Females



* Data are from the National Vital Statistics System (NVSS).

† Rates are age-adjusted to the 2000 U.S. standard population. Rates are suppressed and not ranked if case counts are fewer than 16 or if the population of the specific sex category is less than 50,000.


Note: ONS - Other Nervous System; NOS - Not Otherwise Specified; IBD - Intrahepatic Bile Duct.

Age-Adjusted Invasive Cancer Incidence Rates for the 10 Primary Sites with the Highest Rates within State- and Sex-Specific Categories




New Jersey,^{*†} All Races, 2001

Males



Females



* Data are from selected statewide and metropolitan area cancer registries that met the data quality criteria for publication in *United States Cancer Statistics: 2001 Incidence and Mortality*: case ascertainment is at least 90% complete; $\geq 97\%$ of cases pass a standard set of computerized edits; $\leq 5\%$ of cases were ascertained by death certificate only; $\leq 3\%$ of cases are missing information on sex; $\leq 5\%$ of cases are missing information on race; $\leq 3\%$ of cases are missing information on age; states that are included are shaded in the U.S. map above. U.S. rates cover approximately 92% of the population.

[†] Excludes basal and squamous cell carcinomas of the skin except when these occur on the skin of the genital organs, and *in situ* cancers except urinary bladder.
[‡] Rates are age-adjusted to the 2000 U.S. standard population. Rates are suppressed and not ranked if case counts are fewer than 16 or if the population of the specific sex category is less than 50,000.

Note: ONS - Other Nervous System; NOS - Not Otherwise Specified; IBD - Intrahepatic Bile Duct. Urinary bladder cases include invasive and *in situ*.



Age-Adjusted Cancer Death Rates for the 10 Primary Sites with the Highest Rates within State- and Sex-Specific Categories



New Jersey,* All Races, 2001

Males

Females



* Data are from the National Vital Statistics System (NVSS).

† Rates are age-adjusted to the 2000 U.S. standard population. Rates are suppressed and not ranked if case counts are fewer than 16 or if the population of the specific sex category is less than 50,000.


Note: ONS - Other Nervous System; NOS - Not Otherwise Specified; IBD - Intrahepatic Bile Duct.

Age-Adjusted Invasive Cancer Incidence Rates for the 10 Primary Sites with the Highest Rates within State- and Sex-Specific Categories




New Mexico,^{*†} All Races, 2001

Males



Females



* Data are from selected statewide and metropolitan area cancer registries that met the data quality criteria for publication in *United States Cancer Statistics: 2001 Incidence and Mortality*: case ascertainment is at least 90% complete; $\geq 97\%$ of cases pass a standard set of computerized edits; $\leq 5\%$ of cases were ascertained by death certificate only; $\leq 3\%$ of cases are missing information on sex; $\leq 5\%$ of cases are missing information on race; $\leq 3\%$ of cases are missing information on age; states that are included are shaded in the U.S. map above. U.S. rates cover approximately 92% of the population.

[†] Excludes basal and squamous cell carcinomas of the skin except when these occur on the skin of the genital organs, and *in situ* cancers except urinary bladder.
[‡] Rates are age-adjusted to the 2000 U.S. standard population. Rates are suppressed and not ranked if case counts are fewer than 16 or if the population of the specific sex category is less than 50,000.

Note: ONS - Other Nervous System; NOS - Not Otherwise Specified; IBD - Intrahepatic Bile Duct. Urinary bladder cases include invasive and *in situ*.



Age-Adjusted Cancer Death Rates for the 10 Primary Sites with the Highest Rates within State- and Sex-Specific Categories



New Mexico,* All Races, 2001

Males

Females



* Data are from the National Vital Statistics System (NVSS).

† Rates are age-adjusted to the 2000 U.S. standard population. Rates are suppressed and not ranked if case counts are fewer than 16 or if the population of the specific sex category is less than 50,000.


Note: ONS - Other Nervous System; NOS - Not Otherwise Specified; IBD - Intrahepatic Bile Duct.

Age-Adjusted Invasive Cancer Incidence Rates for the 10 Primary Sites with the Highest Rates within State- and Sex-Specific Categories




New York,^{*†} All Races, 2001

Males



Females



* Data are from selected statewide and metropolitan area cancer registries that met the data quality criteria for publication in *United States Cancer Statistics: 2001 Incidence and Mortality*: case ascertainment is at least 90% complete; $\geq 97\%$ of cases pass a standard set of computerized edits; $\leq 5\%$ of cases were ascertained by death certificate only; $\leq 3\%$ of cases are missing information on sex; $\leq 5\%$ of cases are missing information on race; $\leq 3\%$ of cases are missing information on age; states that are included are shaded in the U.S. map above. U.S. rates cover approximately 92% of the population.

[†] Excludes basal and squamous cell carcinomas of the skin except when these occur on the skin of the genital organs, and *in situ* cancers except urinary bladder.
[‡] Rates are age-adjusted to the 2000 U.S. standard population. Rates are suppressed and not ranked if case counts are fewer than 16 or if the population of the specific sex category is less than 50,000.


Note: ONS - Other Nervous System; NOS - Not Otherwise Specified; IBD - Intrahepatic Bile Duct. Urinary bladder cases include invasive and *in situ*.

Age-Adjusted Cancer Death Rates for the 10 Primary Sites with the Highest Rates within State- and Sex-Specific Categories




New York,* All Races, 2001

Males



Females



* Data are from the National Vital Statistics System (NVSS).

† Rates are age-adjusted to the 2000 U.S. standard population. Rates are suppressed and not ranked if case counts are fewer than 16 or if the population of the specific sex category is less than 50,000.

Note: ONS - Other Nervous System; NOS - Not Otherwise Specified; IBD - Intrahepatic Bile Duct.



Age-Adjusted Invasive Cancer Incidence Rates for the 10 Primary Sites with the Highest Rates within State- and Sex-Specific Categories



North Carolina,^{*†} All Races, 2001

Males

Females



* Data are from selected statewide and metropolitan area cancer registries that met the data quality criteria for publication in *United States Cancer Statistics: 2001 Incidence and Mortality*: case ascertainment is at least 90% complete; $\geq 97\%$ of cases pass a standard set of computerized edits; $\leq 5\%$ of cases were ascertained by death certificate only; $\leq 3\%$ of cases are missing information on sex; $\leq 5\%$ of cases are missing information on race; $\leq 3\%$ of cases are missing information on age; states that are included are shaded in the U.S. map above. U.S. rates cover approximately 92% of the population.

[†] Excludes basal and squamous cell carcinomas of the skin except when these occur on the skin of the genital organs, and *in situ* cancers except urinary bladder.
[‡] Rates are age-adjusted to the 2000 U.S. standard population. Rates are suppressed and not ranked if case counts are fewer than 16 or if the population of the specific sex category is less than 50,000.

Note: ONS - Other Nervous System; NOS - Not Otherwise Specified; IBD - Intrahepatic Bile Duct. Urinary bladder cases include invasive and *in situ*.



Age-Adjusted Cancer Death Rates for the 10 Primary Sites with the Highest Rates within State- and Sex-Specific Categories



North Carolina,* All Races, 2001

Males

Females



* Data are from the National Vital Statistics System (NVSS).

† Rates are age-adjusted to the 2000 U.S. standard population. Rates are suppressed and not ranked if case counts are fewer than 16 or if the population of the specific sex category is less than 50,000.


Note: ONS - Other Nervous System; NOS - Not Otherwise Specified; IBD - Intrahepatic Bile Duct.

Age-Adjusted Cancer Death Rates for the 10 Primary Sites with the Highest Rates within State- and Sex-Specific Categories




North Dakota,* All Races, 2001

Males



Females



* Data are from the National Vital Statistics System (NVSS).

† Rates are age-adjusted to the 2000 U.S. standard population. Rates are suppressed and not ranked if case counts are fewer than 16 or if the population of the specific sex category is less than 50,000.


Note: ONS - Other Nervous System; NOS - Not Otherwise Specified; IBD - Intrahepatic Bile Duct.

Age-Adjusted Invasive Cancer Incidence Rates for the 10 Primary Sites with the Highest Rates within State- and Sex-Specific Categories




Ohio,^{*†} All Races, 2001

Males



Females



* Data are from selected statewide and metropolitan area cancer registries that met the data quality criteria for publication in *United States Cancer Statistics: 2001 Incidence and Mortality*: case ascertainment is at least 90% complete; $\geq 97\%$ of cases pass a standard set of computerized edits; $\leq 5\%$ of cases were ascertained by death certificate only; $\leq 3\%$ of cases are missing information on sex; $\leq 5\%$ of cases are missing information on race; $\leq 3\%$ of cases are missing information on age; states that are included are shaded in the U.S. map above. U.S. rates cover approximately 92% of the population.

[†] Excludes basal and squamous cell carcinomas of the skin except when these occur on the skin of the genital organs, and *in situ* cancers except urinary bladder.
[‡] Rates are age-adjusted to the 2000 U.S. standard population. Rates are suppressed and not ranked if case counts are fewer than 16 or if the population of the specific sex category is less than 50,000.

Note: ONS - Other Nervous System; NOS - Not Otherwise Specified; IBD - Intrahepatic Bile Duct. Urinary bladder cases include invasive and *in situ*.



Age-Adjusted Cancer Death Rates for the 10 Primary Sites with the Highest Rates within State- and Sex-Specific Categories



Ohio,* All Races, 2001

Males

Females



* Data are from the National Vital Statistics System (NVSS).

† Rates are age-adjusted to the 2000 U.S. standard population. Rates are suppressed and not ranked if case counts are fewer than 16 or if the population of the specific sex category is less than 50,000.


Note: ONS - Other Nervous System; NOS - Not Otherwise Specified; IBD - Intrahepatic Bile Duct.

Age-Adjusted Invasive Cancer Incidence Rates for the 10 Primary Sites with the Highest Rates within State- and Sex-Specific Categories




Oklahoma,^{*†} All Races, 2001

Males



Females



* Data are from selected statewide and metropolitan area cancer registries that met the data quality criteria for publication in *United States Cancer Statistics: 2001 Incidence and Mortality*: case ascertainment is at least 90% complete; $\geq 97\%$ of cases pass a standard set of computerized edits; $\leq 5\%$ of cases were ascertained by death certificate only; $\leq 3\%$ of cases are missing information on sex; $\leq 5\%$ of cases are missing information on race; $\leq 3\%$ of cases are missing information on age; states that are included are shaded in the U.S. map above. U.S. rates cover approximately 92% of the population.

[†] Excludes basal and squamous cell carcinomas of the skin except when these occur on the skin of the genital organs, and *in situ* cancers except urinary bladder.
[‡] Rates are age-adjusted to the 2000 U.S. standard population. Rates are suppressed and not ranked if case counts are fewer than 16 or if the population of the specific sex category is less than 50,000.

Note: ONS - Other Nervous System; NOS - Not Otherwise Specified; IBD - Intrahepatic Bile Duct. Urinary bladder cases include invasive and *in situ*.



Age-Adjusted Cancer Death Rates for the 10 Primary Sites with the Highest Rates within State- and Sex-Specific Categories



Oklahoma,* All Races, 2001

Males

Females



* Data are from the National Vital Statistics System (NVSS).

† Rates are age-adjusted to the 2000 U.S. standard population. Rates are suppressed and not ranked if case counts are fewer than 16 or if the population of the specific sex category is less than 50,000.


Note: ONS - Other Nervous System; NOS - Not Otherwise Specified; IBD - Intrahepatic Bile Duct.

Age-Adjusted Invasive Cancer Incidence Rates for the 10 Primary Sites with the Highest Rates within State- and Sex-Specific Categories




Oregon,^{*†} All Races, 2001

Males



Females



* Data are from selected statewide and metropolitan area cancer registries that met the data quality criteria for publication in *United States Cancer Statistics: 2001 Incidence and Mortality*: case ascertainment is at least 90% complete; $\geq 97\%$ of cases pass a standard set of computerized edits; $\leq 5\%$ of cases were ascertained by death certificate only; $\leq 3\%$ of cases are missing information on sex; $\leq 5\%$ of cases are missing information on race; $\leq 3\%$ of cases are missing information on age; states that are included are shaded in the U.S. map above. U.S. rates cover approximately 92% of the population.

[†] Excludes basal and squamous cell carcinomas of the skin except when these occur on the skin of the genital organs, and *in situ* cancers except urinary bladder.
[‡] Rates are age-adjusted to the 2000 U.S. standard population. Rates are suppressed and not ranked if case counts are fewer than 16 or if the population of the specific sex category is less than 50,000.

Note: ONS - Other Nervous System; NOS - Not Otherwise Specified; IBD - Intrahepatic Bile Duct. Urinary bladder cases include invasive and *in situ*.



Age-Adjusted Cancer Death Rates for the 10 Primary Sites with the Highest Rates within State- and Sex-Specific Categories



Oregon,* All Races, 2001

Males

Females



* Data are from the National Vital Statistics System (NVSS).

† Rates are age-adjusted to the 2000 U.S. standard population. Rates are suppressed and not ranked if case counts are fewer than 16 or if the population of the specific sex category is less than 50,000.


Note: ONS - Other Nervous System; NOS - Not Otherwise Specified; IBD - Intrahepatic Bile Duct.

Age-Adjusted Invasive Cancer Incidence Rates for the 10 Primary Sites with the Highest Rates within State- and Sex-Specific Categories




Pennsylvania,^{*,†} All Races, 2001

Males



Females



* Data are from selected statewide and metropolitan area cancer registries that met the data quality criteria for publication in *United States Cancer Statistics: 2001 Incidence and Mortality*: case ascertainment is at least 90% complete; $\geq 97\%$ of cases pass a standard set of computerized edits; $\leq 5\%$ of cases were ascertained by death certificate only; $\leq 3\%$ of cases are missing information on sex; $\leq 5\%$ of cases are missing information on race; $\leq 3\%$ of cases are missing information on age; states that are included are shaded in the U.S. map above. U.S. rates cover approximately 92% of the population.

[†] Excludes basal and squamous cell carcinomas of the skin except when these occur on the skin of the genital organs, and *in situ* cancers except urinary bladder.
[‡] Rates are age-adjusted to the 2000 U.S. standard population. Rates are suppressed and not ranked if case counts are fewer than 16 or if the population of the specific sex category is less than 50,000.


Note: ONS - Other Nervous System; NOS - Not Otherwise Specified; IBD - Intrahepatic Bile Duct. Urinary bladder cases include invasive and *in situ*.

Age-Adjusted Cancer Death Rates for the 10 Primary Sites with the Highest Rates within State- and Sex-Specific Categories




Pennsylvania,* All Races, 2001

Males



Females



* Data are from the National Vital Statistics System (NVSS).

† Rates are age-adjusted to the 2000 U.S. standard population. Rates are suppressed and not ranked if case counts are fewer than 16 or if the population of the specific sex category is less than 50,000.


Note: ONS - Other Nervous System; NOS - Not Otherwise Specified; IBD - Intrahepatic Bile Duct.

Age-Adjusted Invasive Cancer Incidence Rates for the 10 Primary Sites with the Highest Rates within State- and Sex-Specific Categories




Rhode Island,^{*†} All Races, 2001

Males



Females



* Data are from selected statewide and metropolitan area cancer registries that met the data quality criteria for publication in *United States Cancer Statistics: 2001 Incidence and Mortality*: case ascertainment is at least 90% complete; $\geq 97\%$ of cases pass a standard set of computerized edits; $\leq 5\%$ of cases were ascertained by death certificate only; $\leq 3\%$ of cases are missing information on sex; $\leq 5\%$ of cases are missing information on race; $\leq 3\%$ of cases are missing information on age; states that are included are shaded in the U.S. map above. U.S. rates cover approximately 92% of the population.

[†] Excludes basal and squamous cell carcinomas of the skin except when these occur on the skin of the genital organs, and *in situ* cancers except urinary bladder.
[‡] Rates are age-adjusted to the 2000 U.S. standard population. Rates are suppressed and not ranked if case counts are fewer than 16 or if the population of the specific sex category is less than 50,000.


Note: ONS - Other Nervous System; NOS - Not Otherwise Specified; IBD - Intrahepatic Bile Duct. Urinary bladder cases include invasive and *in situ*.

Age-Adjusted Cancer Death Rates for the 10 Primary Sites with the Highest Rates within State- and Sex-Specific Categories




Rhode Island,* All Races, 2001

Males



Females



* Data are from the National Vital Statistics System (NVSS).

† Rates are age-adjusted to the 2000 U.S. standard population. Rates are suppressed and not ranked if case counts are fewer than 16 or if the population of the specific sex category is less than 50,000.

Note: ONS - Other Nervous System; NOS - Not Otherwise Specified; IBD - Intrahepatic Bile Duct.



Age-Adjusted Invasive Cancer Incidence Rates for the 10 Primary Sites with the Highest Rates within State- and Sex-Specific Categories



South Carolina,^{*†} All Races, 2001

Males

Females



* Data are from selected statewide and metropolitan area cancer registries that met the data quality criteria for publication in *United States Cancer Statistics: 2001 Incidence and Mortality*: case ascertainment is at least 90% complete; $\geq 97\%$ of cases pass a standard set of computerized edits; $\leq 5\%$ of cases were ascertained by death certificate only; $\leq 3\%$ of cases are missing information on sex; $\leq 5\%$ of cases are missing information on race; $\leq 3\%$ of cases are missing information on age; states that are included are shaded in the U.S. map above. U.S. rates cover approximately 92% of the population.

[†] Excludes basal and squamous cell carcinomas of the skin except when these occur on the skin of the genital organs, and *in situ* cancers except urinary bladder.
[‡] Rates are age-adjusted to the 2000 U.S. standard population. Rates are suppressed and not ranked if case counts are fewer than 16 or if the population of the specific sex category is less than 50,000.


Note: ONS - Other Nervous System; NOS - Not Otherwise Specified; IBD - Intrahepatic Bile Duct. Urinary bladder cases include invasive and *in situ*.

Age-Adjusted Cancer Death Rates for the 10 Primary Sites with the Highest Rates within State- and Sex-Specific Categories




South Carolina,* All Races, 2001

Males



Females



* Data are from the National Vital Statistics System (NVSS).

† Rates are age-adjusted to the 2000 U.S. standard population. Rates are suppressed and not ranked if case counts are fewer than 16 or if the population of the specific sex category is less than 50,000.

Note: ONS - Other Nervous System; NOS - Not Otherwise Specified; IBD - Intrahepatic Bile Duct.



Age-Adjusted Cancer Death Rates for the 10 Primary Sites with the Highest Rates within State- and Sex-Specific Categories



South Dakota,* All Races, 2001

Males

Females



* Data are from the National Vital Statistics System (NVSS).

† Rates are age-adjusted to the 2000 U.S. standard population. Rates are suppressed and not ranked if case counts are fewer than 16 or if the population of the specific sex category is less than 50,000.

Note: ONS - Other Nervous System; NOS - Not Otherwise Specified; IBD - Intrahepatic Bile Duct.



Age-Adjusted Cancer Death Rates for the 10 Primary Sites with the Highest Rates within State- and Sex-Specific Categories



Tennessee,* All Races, 2001

Males

Females



* Data are from the National Vital Statistics System (NVSS).

† Rates are age-adjusted to the 2000 U.S. standard population. Rates are suppressed and not ranked if case counts are fewer than 16 or if the population of the specific sex category is less than 50,000.


Note: ONS - Other Nervous System; NOS - Not Otherwise Specified; IBD - Intrahepatic Bile Duct.

Age-Adjusted Invasive Cancer Incidence Rates for the 10 Primary Sites with the Highest Rates within State- and Sex-Specific Categories




Texas,^{*†} All Races, 2001

Males



Females



* Data are from selected statewide and metropolitan area cancer registries that met the data quality criteria for publication in *United States Cancer Statistics: 2001 Incidence and Mortality*: case ascertainment is at least 90% complete; $\geq 97\%$ of cases pass a standard set of computerized edits; $\leq 5\%$ of cases were ascertained by death certificate only; $\leq 3\%$ of cases are missing information on sex; $\leq 5\%$ of cases are missing information on race; $\leq 3\%$ of cases are missing information on age; states that are included are shaded in the U.S. map above. U.S. rates cover approximately 92% of the population.

[†] Excludes basal and squamous cell carcinomas of the skin except when these occur on the skin of the genital organs, and *in situ* cancers except urinary bladder.
[‡] Rates are age-adjusted to the 2000 U.S. standard population. Rates are suppressed and not ranked if case counts are fewer than 16 or if the population of the specific sex category is less than 50,000.

Note: ONS - Other Nervous System; NOS - Not Otherwise Specified; IBD - Intrahepatic Bile Duct. Urinary bladder cases include invasive and *in situ*.



Age-Adjusted Cancer Death Rates for the 10 Primary Sites with the Highest Rates within State- and Sex-Specific Categories



Texas,* All Races, 2001

Males

Females



* Data are from the National Vital Statistics System (NVSS).

† Rates are age-adjusted to the 2000 U.S. standard population. Rates are suppressed and not ranked if case counts are fewer than 16 or if the population of the specific sex category is less than 50,000.


Note: ONS - Other Nervous System; NOS - Not Otherwise Specified; IBD - Intrahepatic Bile Duct.

Age-Adjusted Invasive Cancer Incidence Rates for the 10 Primary Sites with the Highest Rates within State- and Sex-Specific Categories




Utah,^{*†} All Races, 2001

Males



Females



* Data are from selected statewide and metropolitan area cancer registries that met the data quality criteria for publication in *United States Cancer Statistics: 2001 Incidence and Mortality*: case ascertainment is at least 90% complete; $\geq 97\%$ of cases pass a standard set of computerized edits; $\leq 5\%$ of cases were ascertained by death certificate only; $\leq 3\%$ of cases are missing information on sex; $\leq 5\%$ of cases are missing information on race; $\leq 3\%$ of cases are missing information on age; states that are included are shaded in the U.S. map above. U.S. rates cover approximately 92% of the population.

[†] Excludes basal and squamous cell carcinomas of the skin except when these occur on the skin of the genital organs, and *in situ* cancers except urinary bladder.
[‡] Rates are age-adjusted to the 2000 U.S. standard population. Rates are suppressed and not ranked if case counts are fewer than 16 or if the population of the specific sex category is less than 50,000.


Note: ONS - Other Nervous System; NOS - Not Otherwise Specified; IBD - Intrahepatic Bile Duct. Urinary bladder cases include invasive and *in situ*.

Age-Adjusted Cancer Death Rates for the 10 Primary Sites with the Highest Rates within State- and Sex-Specific Categories




Utah,* All Races, 2001

Males



Females



* Data are from the National Vital Statistics System (NVSS).

† Rates are age-adjusted to the 2000 U.S. standard population. Rates are suppressed and not ranked if case counts are fewer than 16 or if the population of the specific sex category is less than 50,000.

Note: ONS - Other Nervous System; NOS - Not Otherwise Specified; IBD - Intrahepatic Bile Duct.



Age-Adjusted Invasive Cancer Incidence Rates for the 10 Primary Sites with the Highest Rates within State- and Sex-Specific Categories



Vermont,^{*†} All Races, 2001

Males

Females



* Data are from selected statewide and metropolitan area cancer registries that met the data quality criteria for publication in *United States Cancer Statistics: 2001 Incidence and Mortality*: case ascertainment is at least 90% complete; $\geq 97\%$ of cases pass a standard set of computerized edits; $\leq 5\%$ of cases were ascertained by death certificate only; $\leq 3\%$ of cases are missing information on sex; $\leq 5\%$ of cases are missing information on race; $\leq 3\%$ of cases are missing information on age; states that are included are shaded in the U.S. map above. U.S. rates cover approximately 92% of the population.

[†] Excludes basal and squamous cell carcinomas of the skin except when these occur on the skin of the genital organs, and *in situ* cancers except urinary bladder.
[‡] Rates are age-adjusted to the 2000 U.S. standard population. Rates are suppressed and not ranked if case counts are fewer than 16 or if the population of the specific sex category is less than 50,000.


Note: ONS - Other Nervous System; NOS - Not Otherwise Specified; IBD - Intrahepatic Bile Duct. Urinary bladder cases include invasive and *in situ*.

Age-Adjusted Cancer Death Rates for the 10 Primary Sites with the Highest Rates within State- and Sex-Specific Categories




Vermont,* All Races, 2001

Males



Females



* Data are from the National Vital Statistics System (NVSS).

† Rates are age-adjusted to the 2000 U.S. standard population. Rates are suppressed and not ranked if case counts are fewer than 16 or if the population of the specific sex category is less than 50,000.


Note: ONS - Other Nervous System; NOS - Not Otherwise Specified; IBD - Intrahepatic Bile Duct.

Age-Adjusted Cancer Death Rates for the 10 Primary Sites with the Highest Rates within State- and Sex-Specific Categories




Virginia,* All Races, 2001

Males



Females



* Data are from the National Vital Statistics System (NVSS).

† Rates are age-adjusted to the 2000 U.S. standard population. Rates are suppressed and not ranked if case counts are fewer than 16 or if the population of the specific sex category is less than 50,000.


Note: ONS - Other Nervous System; NOS - Not Otherwise Specified; IBD - Intrahepatic Bile Duct.

Age-Adjusted Invasive Cancer Incidence Rates for the 10 Primary Sites with the Highest Rates within State- and Sex-Specific Categories




Washington,^{*†} All Races, 2001

Males



Females



* Data are from selected statewide and metropolitan area cancer registries that met the data quality criteria for publication in *United States Cancer Statistics: 2001 Incidence and Mortality*: case ascertainment is at least 90% complete; $\geq 97\%$ of cases pass a standard set of computerized edits; $\leq 5\%$ of cases were ascertained by death certificate only; $\leq 3\%$ of cases are missing information on sex; $\leq 5\%$ of cases are missing information on race; $\leq 3\%$ of cases are missing information on age; states that are included are shaded in the U.S. map above. U.S. rates cover approximately 92% of the population.

[†] Excludes basal and squamous cell carcinomas of the skin except when these occur on the skin of the genital organs, and *in situ* cancers except urinary bladder.
[‡] Rates are age-adjusted to the 2000 U.S. standard population. Rates are suppressed and not ranked if case counts are fewer than 16 or if the population of the specific sex category is less than 50,000.


Note: ONS - Other Nervous System; NOS - Not Otherwise Specified; IBD - Intrahepatic Bile Duct. Urinary bladder cases include invasive and *in situ*.

Age-Adjusted Cancer Death Rates for the 10 Primary Sites with the Highest Rates within State- and Sex-Specific Categories




Washington,* All Races, 2001

Males



Females



* Data are from the National Vital Statistics System (NVSS).

† Rates are age-adjusted to the 2000 U.S. standard population. Rates are suppressed and not ranked if case counts are fewer than 16 or if the population of the specific sex category is less than 50,000.

Note: ONS - Other Nervous System; NOS - Not Otherwise Specified; IBD - Intrahepatic Bile Duct.



Age-Adjusted Invasive Cancer Incidence Rates for the 10 Primary Sites with the Highest Rates within State- and Sex-Specific Categories



West Virginia,^{*†} All Races, 2001

Males

Females



* Data are from selected statewide and metropolitan area cancer registries that met the data quality criteria for publication in *United States Cancer Statistics: 2001 Incidence and Mortality*: case ascertainment is at least 90% complete; $\geq 97\%$ of cases pass a standard set of computerized edits; $\leq 5\%$ of cases were ascertained by death certificate only; $\leq 3\%$ of cases are missing information on sex; $\leq 5\%$ of cases are missing information on race; $\leq 3\%$ of cases are missing information on age; states that are included are shaded in the U.S. map above. U.S. rates cover approximately 92% of the population.

[†] Excludes basal and squamous cell carcinomas of the skin except when these occur on the skin of the genital organs, and *in situ* cancers except urinary bladder.
[‡] Rates are age-adjusted to the 2000 U.S. standard population. Rates are suppressed and not ranked if case counts are fewer than 16 or if the population of the specific sex category is less than 50,000.


Note: ONS - Other Nervous System; NOS - Not Otherwise Specified; IBD - Intrahepatic Bile Duct. Urinary bladder cases include invasive and *in situ*.

Age-Adjusted Cancer Death Rates for the 10 Primary Sites with the Highest Rates within State- and Sex-Specific Categories




West Virginia,* All Races, 2001

Males



Females



* Data are from the National Vital Statistics System (NVSS).

† Rates are age-adjusted to the 2000 U.S. standard population. Rates are suppressed and not ranked if case counts are fewer than 16 or if the population of the specific sex category is less than 50,000.

Note: ONS - Other Nervous System; NOS - Not Otherwise Specified; IBD - Intrahepatic Bile Duct.



Age-Adjusted Invasive Cancer Incidence Rates for the 10 Primary Sites with the Highest Rates within State- and Sex-Specific Categories



Wisconsin,^{*†} All Races, 2001

Males

Females



* Data are from selected statewide and metropolitan area cancer registries that met the data quality criteria for publication in *United States Cancer Statistics: 2001 Incidence and Mortality*: case ascertainment is at least 90% complete; $\geq 97\%$ of cases pass a standard set of computerized edits; $\leq 5\%$ of cases were ascertained by death certificate only; $\leq 3\%$ of cases are missing information on sex; $\leq 5\%$ of cases are missing information on race; $\leq 3\%$ of cases are missing information on age; states that are included are shaded in the U.S. map above. U.S. rates cover approximately 92% of the population.

[†] Excludes basal and squamous cell carcinomas of the skin except when these occur on the skin of the genital organs, and *in situ* cancers except urinary bladder.
[‡] Rates are age-adjusted to the 2000 U.S. standard population. Rates are suppressed and not ranked if case counts are fewer than 16 or if the population of the specific sex category is less than 50,000.

Note: ONS - Other Nervous System; NOS - Not Otherwise Specified; IBD - Intrahepatic Bile Duct. Urinary bladder cases include invasive and *in situ*.



Age-Adjusted Cancer Death Rates for the 10 Primary Sites with the Highest Rates within State- and Sex-Specific Categories



Wisconsin,* All Races, 2001

Males

Females



* Data are from the National Vital Statistics System (NVSS).

† Rates are age-adjusted to the 2000 U.S. standard population. Rates are suppressed and not ranked if case counts are fewer than 16 or if the population of the specific sex category is less than 50,000.


Note: ONS - Other Nervous System; NOS - Not Otherwise Specified; IBD - Intrahepatic Bile Duct.

Age-Adjusted Invasive Cancer Incidence Rates for the 10 Primary Sites with the Highest Rates within State- and Sex-Specific Categories




Wyoming,^{*†} All Races, 2001

Males



Females



* Data are from selected statewide and metropolitan area cancer registries that met the data quality criteria for publication in *United States Cancer Statistics: 2001 Incidence and Mortality*: case ascertainment is at least 90% complete; $\geq 97\%$ of cases pass a standard set of computerized edits; $\leq 5\%$ of cases were ascertained by death certificate only; $\leq 3\%$ of cases are missing information on sex; $\leq 5\%$ of cases are missing information on race; $\leq 3\%$ of cases are missing information on age; states that are included are shaded in the U.S. map above. U.S. rates cover approximately 92% of the population.



[†] Excludes basal and squamous cell carcinomas of the skin except when these occur on the skin of the genital organs, and *in situ* cancers except urinary bladder.
[‡] Rates are age-adjusted to the 2000 U.S. standard population. Rates are suppressed and not ranked if case counts are fewer than 16 or if the population of the specific sex category is less than 50,000.

Note: ONS - Other Nervous System; NOS - Not Otherwise Specified; IBD - Intrahepatic Bile Duct. Urinary bladder cases include invasive and *in situ*.

Age-Adjusted Cancer Death Rates for the 10 Primary Sites with the Highest Rates within State- and Sex-Specific Categories



Wyoming,* All Races, 2001



* Data are from the National Vital Statistics System (NVSS).

† Rates are age-adjusted to the 2000 U.S. standard population. Rates are suppressed and not ranked if case counts are fewer than 16 or if the population of the specific sex category is less than 50,000.

Note: ONS - Other Nervous System; NOS - Not Otherwise Specified; IBD - Intrahepatic Bile Duct.